

HOMOLOGICAL SELECTIONS AND FIXED-POINT THEOREMS

V. VALOV

ABSTRACT. A homological selection theorem for C -spaces, as well as a finite-dimensional homological selection theorem is established. We apply the finite-dimensional homological selection theorem to obtain fixed-point theorems forusco homologically UV^n set-valued maps.

1. INTRODUCTION

Banach and Cauty [1, Theorem 8] provided a selection theorem for C -spaces, which is a homological version of the Uspenskij's selection theorem [10, Theorem 1.3]. The aim of this paper was to establish a finite-dimensional form of Banach-Cauty theorem, which is the main tool in proving homological analogues of fixed-point theorems forusco maps established in [2], [4] and [7].

All spaces are assumed to be completely regular. Singular homology $H_n(X; G)$, reduced in dimension 0, with a coefficient group G is considered everywhere below. By default, if not explicitly stated otherwise, G is a ring with unit e . Following the notations from [1], for any space X let $S_k(X; G)$ be the group of all singular chains with coefficients from G consisting of singular k -simplexes and $S(X; G)$ denote the singular complex of X , so $S(X; G)$ is the direct sum $\bigoplus_{k=0}^{\infty} S_k(X; G)$. The groups $S_k(X; G)$ are linked via the boundary homomorphisms $\partial : S_k(X; G) \rightarrow S_{k-1}(X; G)$.

If $\sigma : \Delta^k \rightarrow X$ is a singular k -simplex (Δ^k is the standard k -simplex), we denote by $\|\sigma\|$ the *carrier* $\sigma(\Delta^k)$ of σ . Similarly, we put $\|c\| = \bigcup_i \|\sigma_i\|$ for any chain $c \in S_k(X; G)$, where $c = \sum_i g_i \sigma_i$ is the irreducible representation of c .

For an open cover \mathcal{U} of X let $S(X, \mathcal{U}; G)$ stand for the subgroup of $S(X; G)$ generated by singular simplexes σ with $\|\sigma\| \subset U$ for some

2010 *Mathematics Subject Classification.* Primary 54H25, 55M20; Secondary 55M10, 55M15.

Key words and phrases. fixed points, homological UV^n sets, homological selections.

The author was partially supported by NSERC Grant 261914-13.

$U \in \mathcal{U}$. A homomorphism $\varphi : S(X, \mathcal{U}; G) \rightarrow S(Y; G)$ is said to be a *chain morphism* if $\varphi(S_k(X, \mathcal{U}; G)) \subset S_k(Y; G)$ for all $k \geq 0$ and $\partial \circ \varphi = \varphi \circ \partial$. A point $x \in X$ is called a *fixed point* for a chain morphism $\varphi : S(X, \mathcal{U}; G) \rightarrow S(X; G)$ if for any neighborhood V of x in X there exists a chain $c \in S(V; G) \cap S(X, \mathcal{U}; G)$ such that $\|\varphi(c)\| \cap V \neq \emptyset$.

Let $A \subset B$ be two subsets of a space X . We write $A \xrightarrow{H_m} B$ if the embedding $A \hookrightarrow B$ induces a trivial homomorphism $H_m(A; G) \rightarrow H_m(B; G)$.

A set-valued map $\Phi : X \rightarrow 2^Y$ is called *strongly lower semi-continuous* (br., strongly lsc) if for each compact subset $K \subset Y$ the set $\{x \in X : K \subset \Phi(x)\}$ is open in X . For example, every open-graph set-valued map $\Phi : X \rightarrow 2^Y$ is strongly lsc, see [5, Proposition 3.2].

Here is our first homological selection theorem.

Theorem 1.1. *Let X be a paracompact C -space, Y be an arbitrary space and $\Phi_k : X \rightarrow 2^Y$, $m = 0, 1, \dots, n$, be a finite sequence of strongly lsc maps satisfying the following conditions, where G is a fixed ring with unit:*

- (i) $\Phi_m(x) \xrightarrow{H_m} \Phi_{m+1}(x)$ for every $m = 0, \dots, n-1$ and every $x \in X$;
- (ii) $H_m(\Phi_n(x); G) = 0$ for all $m \geq n$ and all $x \in X$.

Then there exists an open cover \mathcal{U} of X and a chain morphism $\varphi : S(X, \mathcal{U}; G) \rightarrow S(Y; G)$ such that $\varphi(S(U; G)) \subset S(\Phi_n(x); G)$ for every $U \in \mathcal{U}$ and every $x \in U$.

Let us mention that the Banach-Cauty result [1, Theorem 8] is a particular case of Theorem 1.1 with $\Phi_m = \Phi_n$ for all m . There is also a finite-dimensional analogue of the above theorem.

Theorem 1.2. *Let X , Y and G be as in Theorem 1.1. The same conclusion holds if $\dim X \leq n$ and the sequence of strongly lsc maps $\Phi_m : X \rightarrow 2^Y$ satisfies only condition (i).*

Theorem 1.2 and [1, Theorem 7] imply the following fixed point theorem for usco (upper semi-continuous and compact-valued) maps:

Theorem 1.3. *Let X be a paracompact space with $\dim X \leq n$, Y a compact metric AR, G a field and $\Phi : X \rightarrow 2^Y$ be a homologically $UV^{n-1}(G)$ usco map. Then for every continuous map $g : Y \rightarrow X$ there exists a point $y_0 \in Y$ with $y_0 \in \Phi(g(y_0))$.*

The particular case of Theorem 1.3 with $X = Y$ and $g = id_X$ is also interesting.

Corollary 1.4. *Let X be a compact metric AR with $\dim X \leq n$, G a field and $\Phi : X \rightarrow 2^X$ be a homologically $UV^{n-1}(G)$ usco map. Then there exists a point $x_0 \in X$ with $x_0 \in \Phi(x_0)$.*

Recall that a closed subset A of a metric space X is called UV^n in X if every neighborhood U of A in X contains another neighborhood V such that the inclusion $V \hookrightarrow U$ generates trivial homomorphisms $\pi_k(V) \rightarrow \pi_k(U)$ between the homotopy groups for all $k = 0, \dots, n$. If considering the homology groups $H_k(\cdot; G)$ instead of the homotopy groups $\pi_k(V)$ and $\pi_k(U)$ (i.e. requiring $V \xrightarrow{H_m} U$ for all $m = 0, 1, \dots, n$), then A is said to be *homologically $UV^n(G)$ in X* . It follows from the universal formula coefficients that every UV^n -subset of X is homologically $UV^n(G)$ in X for all groups G . Moreover, following the proof of Proposition 7.1.3 from [8], one can show that A is homologically $UV^n(G)$ in a given metric ANR -space X if and only if it is homologically $UV^n(G)$ in any metric ANR -space that contains A as a closed set. We say that A is *homologically $UV^\omega(G)$ in X* if A is homologically $UV^n(G)$ for all $n \geq 0$. Because of the last two notions, it is convenient to say that A is *homotopically UV^n in X* instead of A being UV^n in X . We also say that a set-valued map $\Phi : X \rightarrow 2^Y$ is *homologically $UV^n(G)$* if all values $\Phi(x)$ are homologically $UV^n(G)$ -subsets of Y .

Theorem 1.5. *Let X be a compact metric AR -space, G a field and $\Phi : X \rightarrow 2^X$ be a homologically $UV^\omega(G)$ usco map. Then there exists a point $x_0 \in Y$ with $x_0 \in \Phi(x_0)$.*

Theorem 1.3 was established by Gutev [7] for usco maps with homotopically UV^n values. A homotopical version of Theorem 1.5 is also known, see Corollaries 3.6 and 5.14 from [4], or Theorem 1.3 from [7]. One can also show that if X is a compact metric AR and $\Phi : X \rightarrow 2^X$ is a homologically $UV^\omega(G)$ usco map, then each value $\Phi(x)$ has trivial Čech homology groups with coefficients in G . So, in the particular case when G is the group \mathbb{Q} of the rationals, Theorem 1.5 follows from the more general [3, Theorem 7] treating the so-called algebraic AR 's. However, in the framework of usual AR 's Theorem 1.5 provides a very simple proof.

2. HOMOLOGICAL SELECTION THEOREMS

In this section we prove Theorems 1.1 - 1.2. For any simplicial complex K and an integer $m \geq 0$ let $K^{(m)}$ and $C_m(K; G)$ denote, respectively, the m -skeleton of K and the group generated by the oriented m -simplexes of K with coefficients in G .

We say that a chain morphism $\mu : C(K; G) \rightarrow S(A; G)$ (resp., $\mu : S(A; G) \rightarrow C(K; G)$), where K is a simplicial complex and A a topological space, is *correct* provided $\mu(v)$ is a singular 0-simplex in

$S(A; G)$ for every vertex $v \in K^{(0)}$ (resp., $\mu(\sigma)$ is a vertex of K for every singular 0-simplex $\sigma \in S_0(A; G)$).

Lemma 2.1. *Suppose $\{A\}_{k=0}^{m+1}$ is a sequence of subsets of Y with $A_k \xrightarrow{H_k} A_{k+1}$, $k = 0, 1, \dots, m$. Let L be a simplicial complex of dimension m and K be the cone of L . Then every correct chain morphism $\mu_m : C(L; G) \rightarrow S_m(A_m; G)$ such that $\mu_m(C(L^{(k)}; G)) \subset S_k(A_k; G)$ for all $k \leq m$ can be extended to a correct chain morphism $\mu_{m+1} : C(K; G) \rightarrow S_{m+1}(A_{m+1}; G)$ satisfying the following conditions:*

- $\mu_{m+1}(C(K^{(k)}; G)) \subset S_k(A_k; G)$ for all $k = 0, 1, \dots, m+1$;
- $\tilde{\mu}_m \circ \partial_{m+1} = \partial_{m+1} \circ \mu_{m+1}$, where $\tilde{\mu}_m = \mu_{m+1}|_{C(K^{(m)}; G)}$.

Proof. We first extend each morphism $\mu_k = \mu_m|_{C(L^{(k)}; G)}$ to a morphism $\tilde{\mu}_k : C(K^{(k)}; G) \rightarrow S_k(A_k; G)$ such that $\tilde{\mu}_k \circ \partial_{k+1} = \partial_{k+1} \circ \tilde{\mu}_{k+1}$, $k = 0, 1, \dots, m-1$. To this end, denote by v_0 the vertex of K and consider the augmentation $\epsilon : S_0(A_1; G) \rightarrow G$ defined by $\epsilon(\sigma) = e$ for all singular 0-simplexes $\sigma \in S_0(A_1; G)$. Define $\tilde{\mu}_0(\{v_0\})$ to be a fixed singular simplex $\sigma_0 \in S_0(A_0; G)$ and $\tilde{\mu}_0(\{v\}) = \mu_0(\{v\})$ for any $v \in L^{(0)}$. Then extend μ_0 to a homomorphism $\tilde{\mu}_0 : C(K^{(0)}; G) \rightarrow S_0(A_0; G)$ by linearity.

If $\sigma = (v_1, v_2)$ is an 1-dimensional simplex in K , then $\tilde{\mu}_0(\partial_1(\sigma)) = \tilde{\mu}_0(v_2) - \tilde{\mu}_0(v_1)$. Hence, $\epsilon(\tilde{\mu}_0(\partial_1(\sigma))) = 0$. Since $A_0 \xrightarrow{H_0} A_1$, there is a singular chain $\tau_\sigma \in S_1(A_1; G)$ such that $\tilde{\mu}_0(\partial_1(\sigma)) = \partial_1(\tau_\sigma)$. Letting $\tilde{\mu}_1(\sigma) = \tau_\sigma$ if $\sigma \in K^{(1)} \setminus L^{(1)}$ and $\tilde{\mu}_1(\sigma) = \mu_1(\sigma)$ if $\sigma \in L^{(1)}$, we define the homomorphism $\tilde{\mu}_1$ on every simplex of $K^{(1)}$. Then extend this homomorphism to $\tilde{\mu}_1 : C(K^{(1)}; G) \rightarrow S_1(A_1; G)$ by linearity.

Because $A_{k-1} \xrightarrow{H_{k-1}} A_k$, we can repeat the above construction to obtain the homomorphisms $\tilde{\mu}_k$ for any $k \leq m$. Then $\tilde{\mu}_m : C(K^{(m)}; G) \rightarrow S_m(A_m; G)$. Since $A_m \xrightarrow{H_m} A_{m+1}$, we can use once more the above arguments to obtain the chain morphism $\mu_{m+1} : C(K; G) \rightarrow S_{m+1}(A_{m+1}; G)$ satisfying the required conditions. \square

Lemma 2.2. *Let L be a simplicial complex with trivial homology groups and $A \subset B$ be a pair of spaces. Then every correct chain morphism $\nu : S(A; G) \rightarrow C(L; G)$ can be extended to a correct chain morphism $\tilde{\nu} : S(B; G) \rightarrow C(L; G)$.*

Proof. We are going to define by induction for each $k \geq 0$ a homomorphism $\tilde{\nu}_k : S_k(B; G) \rightarrow C_k(L; G)$ extending $\nu_k : S_k(A; G) \rightarrow C_k(L; G)$ such that $\tilde{\nu}_k(\partial_{k+1}(c)) = \partial_{k+1}(\tilde{\nu}_{k+1}(c))$ for every singular chain $c \in S_{k+1}(B; G)$ and $k \geq 0$. For every singular 0-simplex $\sigma \in S_0(B; G)$ we define $\tilde{\nu}_0(\sigma) = \nu_0(\sigma)$ if $\sigma \in S_0(A; G)$ and $\tilde{\nu}_0(\sigma) = v_0$ if $\sigma \notin S_0(A; G)$,

where v_0 is a fixed vertex of L . Then extend this homomorphism over $S_0(B; G)$ by linearity. Because ν is correct, $\tilde{\nu}_0(\sigma)$ is a vertex of L for all singular 0-simplexes $\sigma \in S(B; G)$.

To define $\tilde{\nu}_1$ we consider the augmentation $\epsilon : C_0(L; G) \rightarrow G$ defined by $\epsilon(v) = e$ for all vertexes of L , see [9]. Thus, $\epsilon(\tilde{\nu}_0(\partial_1(\sigma))) = 0$ for every singular 1-simplex $\sigma \in S_1(B; G)$. Because $H_0(L; G) = 0$, $\partial_1(C_1(L; G)) = \epsilon^{-1}(0)$. Therefore, for every singular simplex $\sigma \in S_1(B; G) \setminus S_1(A; G)$ there exists a chain $c_\sigma \in C_1(L; G)$ such that $\partial_1(c_\sigma) = \tilde{\nu}_0(\partial_1(\sigma))$. We define $\tilde{\nu}_1(\sigma) = \nu_1(\sigma)$ if $\sigma \in S_1(A; G)$ and $\tilde{\nu}_1(\sigma) = c_\sigma$ if $\sigma \in S_1(B; G) \setminus S_1(A; G)$, and extend $\tilde{\nu}_1$ over $S_1(B; G)$ by linearity.

Suppose the homomorphism $\tilde{\nu}_k : S_k(B; G) \rightarrow C_k(L; G)$ was already constructed. Then, using that the kernel of the boundary homomorphism $\partial_k : C_k(L; G) \rightarrow C_{k-1}(L; G)$ coincides with the image $\partial_{k+1}(C_{k+1}(L; G))$, we can define $\tilde{\nu}_{k+1}$ extending $\tilde{\nu}_k$ and satisfying the equality $\tilde{\nu}_k \circ \partial_{k+1} = \partial_{k+1} \circ \tilde{\nu}_{k+1}$. \square

Proof of Theorem 1.1. We modify the proof of [1, Theorem 8]. By induction we are going to construct two sequences of locally finite open covers of X , $\mathcal{V}_m = \{V_\alpha : \alpha \in \Gamma_m\}$ and $\mathcal{W}_m = \{W_\alpha : \alpha \in \Gamma_m\}$, $m \geq 0$, an increasing sequence $K_0 \subset K_1 \subset \dots$ of simplicial complexes and correct chain morphisms $\mu_m : C(K_m; G) \rightarrow S_m(Y; G)$ such that

- (1) $\overline{W}_\alpha \subset V_\alpha$ for all $\alpha \in \Gamma_m$, $m \geq 0$;
- (2) $\dim K_m = m$;
- (3) $\mu_{m+1}|_{C(K_m; G)} = \mu_m$ and $\partial_{m+1} \circ \mu_{m+1} = \tilde{\mu}_m \circ \partial_{m+1}$, where $\tilde{\mu}_m = \mu_{m+1}|_{C(K_m^{(m)}; G)}$.

Moreover, for every m and $\alpha \in \Gamma_m$ we shall assign a finite sub-complex L_α of K_m and a set $\Omega_\alpha = \bigcup_{\sigma \in L_\alpha} \|\mu_m(\sigma)\|$ satisfying the following conditions:

- (4) $\dim L_\alpha = m$ and L_α is a cone whose base is a sub-complex $M_\alpha \subset K_{m-1}$ and having a vertex α ;
- (5) If $m \leq n$ and $\alpha \in \Gamma_m$, then $\Omega_\alpha \subset \Phi_m(x)$ and $\Omega_\alpha^{(k)} \subset \Phi_k(x)$ for all $k \leq m-1$ and all $x \in V_\alpha$, where $\Omega_\alpha^{(k)} = \bigcup_{\sigma \in L_\alpha} \|\mu_m(\sigma^{(k)})\|$;
- (6) If $m > n$ and $\alpha \in \Gamma_m$, then $\Omega_\alpha \subset \Phi_n(x)$ for all $x \in V_\alpha$.

To start our construction, for every $x \in X$ we fix a point $y_x \in \Phi_0(x)$ and consider the set $O_x = \{x' \in X : y_x \in \Phi_0(x')\}$. Since Φ_0 is strongly lsc, O_x is open in X . Let $\mathcal{V}_0 = \{V_\alpha : \alpha \in \Gamma_0\}$ be a locally finite open cover of X refining the cover $\{O_x : x \in X\}$, and choose $\mathcal{W}_0 = \{W_\alpha : \alpha \in \Gamma_0\}$ to be a locally finite open cover of X with $\overline{W}_\alpha \subset V_\alpha$ for all $\alpha \in \Gamma_0$. Let the complex K_0 be the zero-dimensional complex whose set of vertices is Γ_0 . For every $\alpha \in \Gamma_0$ we set $L_\alpha = \{\alpha\}$ and choose $x_\alpha \in X$ such that $V_\alpha \subset O_{x_\alpha}$. Define $\mu_0 : C(K_0; G) \rightarrow S_0(Y; G)$ to be

the homomorphism assigning to each generator corresponding to α the singular 0-simplex y_{x_α} , and let $\Omega_\alpha = \{y_{x_\alpha}\}$. Obviously, μ_0 is correct.

Suppose for some $m < n - 1$ and all $k \leq m$ we already performed the construction satisfying conditions (1) – (5). Then for every $x \in X$ choose an open neighborhood G_x of x meeting only finitely many elements of the cover $\bigcup_{k \leq m} \mathcal{V}_k$ such that $G_x \subset V_\alpha$ for all $\alpha \in \bigcup_{k=0}^m \Gamma_k$ with $G_x \cap \overline{W}_\alpha \neq \emptyset$. Let $J(x) = \{\alpha \in \bigcup_{k=0}^m \Gamma_k : G_x \subset V_\alpha\}$ and $D_x^{(k)} = \bigcup \{\Omega_\alpha^{(k)} : \alpha \in J(x)\}$, $k \leq m$. Since $J(x)$ is finite, all $D_x^{(k)}$ are compact subsets of Y with $D_x^{(k)} \subset D_x^{(m)} = D_x$. Moreover, condition (5) implies $D_x \subset \Phi_m(x)$. Consider the finite sub-complex $M_x = \bigcup \{L_\alpha : \alpha \in J(x)\}$ of K_m and the cone L_x with a vertex $v_x \notin K_m$ and a base M_x . Then, according to the definition of Ω_α and condition (5), we have $\mu_m(C(M_x^{(k)}; G)) \subset S_k(D_x^{(k)}; G) \subset S_k(\Phi_k(x); G)$, $k \leq m$. Therefore, we can apply Lemma 2.1 to find a correct chain morphism $\mu_x : C(L_x; G) \rightarrow S_{m+1}(\Phi_{m+1}(x); G)$ extending $\mu_m|C(M_x; G)$ such that $\mu_x(C(L_x^{(k)}; G)) \rightarrow S_k(\Phi_k(x); G)$ and $\partial_{m+1} \circ \mu_x = (\mu_x|C(L_x^{(m)}; G)) \circ \partial_x$, where $\partial_x : C(L_x; G) \rightarrow C(L_x^{(m)}; G)$ is the boundary homomorphism. Then $\Omega_x = \bigcup_{\sigma \in L_x} \|\mu_x(\sigma)\|$ is a compact subset of $\Phi_{m+1}(x)$ containing D_x . The strong lower semi-continuity of Φ_{m+1} yields that $O_x^m = \{x' \in G_x : \Omega_x \subset \Phi_{m+1}(x')\}$ is an open neighborhood of x . So, there exists a locally finite open cover $\mathcal{V}_{m+1} = \{V_\alpha : \alpha \in \Gamma_{m+1}\}$ of X refining the cover $\{O_x^m : x \in X\}$, and take a locally finite open cover $\mathcal{W}_{m+1} = \{W_\alpha : \alpha \in \Gamma_{m+1}\}$ satisfying condition (1). Now, for every $\alpha \in \Gamma_{m+1}$ choose $x_\alpha \in X$ with $V_\alpha \subset O_{x_\alpha}^m$ and let L_α be the cone with base M_{x_α} and vertex α . Define K_{m+1} to be the union $K_m \cup \bigcup_{\alpha \in \Gamma_{m+1}} L_\alpha$. Identifying the cones L_α and L_{x_α} , we define the correct morphism $\mu_{m+1} : C(K_{m+1}; G) \rightarrow S_{m+1}(Y; G)$ by $\mu_{m+1}|C(K_m; G) = \mu_m$ and $\mu_{m+1}|C(L_\alpha; G) = \mu_{x_\alpha}$. Finally, let $\Omega_\alpha = \Omega_{x_\alpha}$. It is easily seen that conditions (1) – (5) are satisfied. Moreover, the definition of G_x and the inclusion $O_{x_\alpha}^m \subset G_{x_\alpha}$ yield that

- (7) For every $\beta \in \bigcup_{k=0}^m \Gamma_k$ and $\alpha \in \Gamma_{m+1}$ with $W_\alpha \cap W_\beta \neq \emptyset$ we have $W_\alpha \subset V_\beta$, and thus $L_\beta \subset L_\alpha$.

In this way we can perform our construction for all $m \leq n$. If we substitute $\Phi_m = \Phi_n$ for all $m \geq n$, we have also $\Phi_m(x) \xrightarrow{H_m} \Phi_{m+1}(x)$ because $H_m(\Phi_n(x); G) = 0$ for all $x \in X$. Therefore, following the above arguments, we can perform for all m the steps from m to $m + 1$ satisfying conditions (1) – (7).

Let $K = \bigcup_{m=0}^\infty K_m$. Then the morphisms μ_m define a correct chain morphism $\mu : C(K; G) \rightarrow S(Y; G)$. Because X is a C -space, there

exists a sequence of disjoint open families $\mathcal{U}_m = \{U_\lambda : \lambda \in \Lambda_m\}$, $m \geq 0$, such that each \mathcal{U}_m refines \mathcal{W}_m and the family $\mathcal{U} = \bigcup_{m=0}^{\infty} \mathcal{U}_m$ covers X .

The final step of the proof is to construct of a chain morphism from $S(X, \mathcal{U}; G)$ into $C(K; G)$. To this end, let $\Lambda = \bigcup_{k=0}^{\infty} \Lambda_k$ and $\Lambda(m) = \bigcup_{k=0}^m \Lambda_k$. Consider also the sub-complexes $S_m = \sum_{\lambda \in \Lambda(m)} S(U_\lambda; G)$ of $S(X; G)$, $m \geq 0$, whose union is $S(X, \mathcal{U}; G)$. For every $\lambda \in \Lambda_m$ select an $\alpha_\lambda \in \Gamma_m$ with $U_\lambda \subset W_{\alpha_\lambda}$. We are going to construct a correct chain morphism $\nu : S(X, \mathcal{U}; G) \rightarrow C(K; G)$ such that

$$(8) \quad \nu(S(U_\lambda; G)) \subset C(L_{\alpha_\lambda}; G) \text{ for all } \lambda \in \Lambda.$$

For any $\lambda \in \Lambda_0$ the complex L_{α_λ} is a single point. So, we can find a chain morphism $\nu_\lambda : S(U_\lambda; G) \rightarrow C(L_{\alpha_\lambda}; G)$. Since the family \mathcal{U}_0 is disjoint, S_0 is the direct sum of all $S(U_\lambda; G)$, $\lambda \in \Lambda_0$. Hence, the chain morphism $\nu_0 : S_0 \rightarrow C(K; G)$ with $\nu_0|_{S(U_\lambda; G)} = \nu_\lambda$ for all $\lambda \in \Lambda_0$ is well defined and $\nu_0(S(U_\lambda; G)) \subset C(L_{\alpha_\lambda}; G)$.

Suppose that for some m we have constructed correct chain morphisms $\nu_k : S_k \rightarrow C(K; G)$, $k \leq m$, such that ν_k extends ν_{k-1} and $\nu_k(S(U_\lambda; G)) \subset C(L_{\alpha_\lambda}; G)$ for all $\lambda \in \Lambda(k)$. Because \mathcal{U}_{m+1} is a disjoint family, so is the family $\{S(U_\lambda; G) : \lambda \in \Lambda_{m+1}\}$. Therefore, to extend ν_m over S_{m+1} , it suffices for every $\lambda \in \Lambda_{m+1}$ to extend $\nu_m|_{(S(U_\lambda; G) \cap S_m)}$ over $S(U_\lambda; G)$. To this end, observe that if $\lambda \in \Lambda_{m+1}$ and $\lambda' \in \Lambda(m)$ with $U_\lambda \cap U_{\lambda'} \neq \emptyset$, then $W_{\alpha_\lambda} \cap W_{\alpha_{\lambda'}} \neq \emptyset$. Thus, according to condition (7), $L_{\alpha_{\lambda'}} \subset L_{\alpha_\lambda}$. Consequently, by (8), $\nu_m(S(U_\lambda; G) \cap S_m) \subset C(L_{\alpha_\lambda}; G)$ for any $\lambda \in \Lambda_{m+1}$. Since L_{α_λ} is contractible and ν_m is correct, we can apply Lemma 2.2 (with $A = U_\lambda \cap \bigcup_{\lambda' \in \Lambda(m)} U_{\lambda'}$ and $B = U_\lambda$) to find a correct chain morphism $\nu_\lambda : S(U_\lambda; G) \rightarrow C(L_{\alpha_\lambda}; G)$ extending $\nu_m|_{S(U_\lambda; G)}$. This completes the induction, so the construction of the required chain morphism $\nu : S(X, \mathcal{U}; G) \rightarrow C(K; G)$ is done.

Finally, let $\varphi : S(X, \mathcal{U}; G) \rightarrow S(Y; G)$ be the composition $\varphi = \mu \circ \nu$. Then, according to (7) and the definitions of Ω_α , for every $\lambda \in \Lambda$ we have

$$\varphi(S(U_\lambda; G)) \subset \mu(C(L_{\alpha_\lambda}; G)) \subset S(\Omega_{\alpha_\lambda}; G).$$

Since $U_\lambda \subset W_{\alpha_\lambda}$, conditions (4) and (5) yield that $\Omega_{\alpha_\lambda} \subset \Phi_n(x)$ for all $x \in U_\lambda$. Therefore, $\varphi(S(U_\lambda; G))$ is contained in $S(\Phi_n(x); G)$ whenever $x \in U_\lambda$. \square

Proof of Theorem 1.2. Since the sequence $\{\Phi_m\}_{m=0}^n$ satisfies condition (i) from Theorem 1.1, we can perform the construction from the proof of Theorem 1.1 for every $m = 0, 1, \dots, n$. So, we construct the locally finite covers $\mathcal{V}_m = \{V_\alpha : \alpha \in \Gamma_m\}$ and $\mathcal{W}_m = \{W_\alpha : \alpha \in \Gamma_m\}$ of X , the complexes $K_0 \subset K_1 \subset \dots \subset K_n$, the sets Ω_α for any $\alpha \in \bigcup_{k=0}^n \Gamma_k$ and the correct chain morphisms $\mu_m : C(K_m; G) \rightarrow S_m(Y; G)$ satisfying conditions (1) – (5) and the particular case of condition (7)

with $m \leq n - 1$. Since the complex $K = \bigcup_{m=0}^n K_m$ is n -dimensional, $K^{(m)} = \emptyset$ for all $m > n$. So, we can suppose that $\mu_m = \mu_n$ for $m \geq n$. In this way we obtain a chain morphism $\mu : C(K; G) \rightarrow S(Y; G)$. Because $\dim X \leq n$, according to Corollary 5.3 from [6], for every $m = 0, 1, \dots, n$ there exists a disjoint family $\mathcal{U}_m = \{U_\lambda : \lambda \in \Lambda_m\}$ such that each \mathcal{U}_m refines \mathcal{W}_m and the family $\mathcal{U} = \bigcup_{m=0}^n \mathcal{U}_m$ covers X . Then, repeating the arguments from the final part of the proof of Theorem 1.1, we construct the required chain morphism $\varphi : S(X, \mathcal{U}; G) \rightarrow S(Y; G)$. \square

3. FIXED-POINT THEOREMS FOR HOMOLOGICALLY $UV^n(G)$ USCO MAPS

In this section we prove Theorems 1.3 and 1.5. For a set-valued map $\Phi : X \rightarrow 2^Y$ we denote by $\mathcal{O}(\Phi)$ the family of the open-graph maps $\Theta : X \rightarrow 2^Y$ such that $\Phi(x) \subset \Theta(x)$ for all $x \in X$. Next proposition is a homological version (and its proof is a small modification) of [5, Proposition 4.2].

Proposition 3.1. *Let X be a paracompact space, Y be a space and let $\Phi : X \rightarrow 2^Y$ be an usco map such that for every $x \in X$ and a neighborhood U of $\Phi(x)$ there exists a neighborhood V of $\Phi(x)$ with $V \xrightarrow{H_m} U$. Then for every $\varphi \in \mathcal{O}(\Phi)$ there exists $\Theta \in \mathcal{O}(\Phi)$ such that $\Theta(x)$ is open in Y and $\overline{\Theta(x)} \xrightarrow{H_m} \varphi(x)$ for all $x \in X$.*

Proof. Let $\varphi \in \mathcal{O}(\Phi)$. Then the graph $G(\varphi)$ is open in $X \times Y$ and contains the compact set $\{x\} \times \Phi(x)$ for every $x \in X$. So, there exist neighborhoods $W_1(x)$ and $U(x)$ of x and $\Phi(x)$, respectively, with $W_1(x) \times U(x) \subset G(\varphi)$. Thus, $\Phi(x) \subset U(x) \subset \varphi(x')$ for all $x' \in W_1(x)$. Then $\overline{V(x)} \xrightarrow{H_m} U(x)$ for some open neighborhood $V(x)$ of $\Phi(x)$. Since Φ is upper semi-continuous, we can find a neighborhood $W(x) \subset W_1(x)$ such that $x' \in W(x)$ implies $\Phi(x') \subset V(x)$. Hence, for all $x' \in W(x)$ we have

$$(9) \quad \Phi(x') \subset \overline{V(x)} \xrightarrow{H_m} U(x) \subset \varphi(x').$$

Next, let $\gamma = \{P_\alpha : \alpha \in A\}$ be a locally finite closed cover of X refining the cover $\{W(x) : x \in X\}$ (recall that X is paracompact), and for every α fix $x_\alpha \in X$ such that $P_\alpha \subset W(x_\alpha)$. For every $x \in X$ the set $A(x) = \{\alpha \in A : x \in P_\alpha\}$ is finite, and define $\Theta(x) = \bigcap \{V(x_\alpha) : \alpha \in A(x)\}$. One can show that Θ is open-graph (see the proof of [5, Proposition 4.2]). Moreover, since $x \in \bigcap \{W(x_\alpha) : \alpha \in A(x)\}$, it follows from (9) that

$$\Phi(x) \subset \overline{V(x_\alpha)} \xrightarrow{H_m} U(x_\alpha) \subset \varphi(x)$$

for all $\alpha \in A(x)$. This yields $\Phi(x) \subset \Theta(x) \subset \overline{\Theta(x)} \xrightarrow{H_m} \varphi(x)$. \square

Proof of Theorem 1.3. Let $g : Y \rightarrow X$ be a continuous (single-valued) map. Without loss of generality, we may assume that $g(Y) = X$. We need to show that the set-valued map $\Phi_g = \Phi \circ g : Y \rightarrow 2^Y$ has a fixed-point. Suppose this is not true. So $y \notin \Phi_g(y)$ for all $y \in Y$, or equivalently $\Phi(x) \subset Y \setminus g^{-1}(x)$ for all $x \in X$. Consider the set-valued map $\varphi : X \rightarrow 2^Y$, $\varphi(x) = Y \setminus g^{-1}(x)$. Then Φ is a selection for φ and it is easily seen that φ has an open graph. Because Φ is homologically $UV^{n-1}(G)$, we can apply Proposition 3.1 to find for each $m = 0, 1, \dots, n$ a set valued map $\Theta_m : X \rightarrow 2^Y$ such that

- $\Phi(x) \subset \Theta_0(x)$, $x \in X$;
- $\Theta_m(x) \xrightarrow{H_m} \Theta_{m+1}(x)$ for all $x \in X$ and $m = 0, \dots, n-1$;
- Each $\Theta_n(x)$ is open in Y and $\overline{\Theta_n(x)} \subset \varphi(x)$, $x \in X$.

Then, according to Theorem 1.2, there exists an open cover \mathcal{U} of X and a chain morphism $\phi : S(X, \mathcal{U}; G) \rightarrow S(Y; G)$ such that $\phi(S(U; G)) \subset S(\Theta_n(x); G)$ for every $U \in \mathcal{U}$ and every $x \in U$. Consider the open cover $\mathcal{U}_g = g^{-1}(\mathcal{U})$ of Y and the chain morphism $g_\# : S(Y, \mathcal{U}_g; G) \rightarrow S(X, \mathcal{U}; G)$ generated by g . Then $\phi_g = \phi \circ g_\# : S(Y, \mathcal{U}_g; G) \rightarrow S(Y; G)$ is a chain morphism with

$$(10) \quad \phi_g(S(g^{-1}(U); G)) \subset S(\Theta_n(g(y)); G) \text{ for all } U \in \mathcal{U} \text{ and } y \in g^{-1}(U).$$

So, we can apply the homological fixed-point theorem [1, Theorem 7] to conclude that the chain morphism ϕ_g has a fixed point $y_0 \in Y$. This means that for any neighborhood $W \subset Y$ of y_0 there is a chain $c_W \in S(W; G) \cap S(Y, \mathcal{U}_g; G)$ such that $\|\phi_g(c_W)\| \cap W \neq \emptyset$. Choose $U_0 \in \mathcal{U}$ with $y_0 \in g^{-1}(U_0)$ and let $V \subset g^{-1}(U_0)$ be a neighborhood of y_0 . Then $c_V \in S(V; G) \subset S(g^{-1}(U_0); G)$. Thus, we have

$$(11) \quad \|\phi_g(c_V)\| \cap V \neq \emptyset \text{ and, by (10), } \|\phi_g(c_V)\| \subset \Theta_n(g(y_0)).$$

On the other hand, since $\overline{\Theta_n(x_0)} \subset \varphi(x_0)$, where $x_0 = g(y_0)$, we can choose V to be so small that $V \cap \overline{\Theta_n(x_0)} = \emptyset$. The last relation contradicts condition (11). \square

Proof of Theorem 1.5. The arguments from the proof of [5, Theorem 1.3] work in our situation. For completeness, we provide a sketch. Since X can be embedded in the Hilbert cube Q as a retract, we may suppose that $\Phi : Q \rightarrow 2^Q$ is a homologically $UV^\omega(G)$ usco map. Identifying Q with the product $\prod \{\mathbb{I}_k : k \in \mathbb{N}\}$, where $\mathbb{I} = [0, 1]$, let $\pi_n : Q \rightarrow \prod \{\mathbb{I}_k : k \leq n\}$ be the projection onto the cube \mathbb{I}^n and $h_n : \mathbb{I}^n \rightarrow Q$ be the embedding assigning to every point $x = (x_1, \dots, x_n) \in \mathbb{I}^n$ the point $h(x)$ having the same first n -coordinates and all other coordinates 0.

For every n consider the homologically $UV^\omega(G)$ usco map $\Phi_n : \mathbb{I}^n \rightarrow 2^Q$ defined by $\Phi_n(x) = \Phi(h_n(x))$. Then, according to Theorem 1.3 (with $X = \mathbb{I}^n$, $Y = Q$, $g = \pi_n$ and $\Phi = \Phi_n$), there is a point $x^n \in Q$ with $x^n \in \Phi_n(\pi_n(x^n))$. If $x^0 \in Q$ is the limit point of a convergent subsequence of $\{x^n\}_{n \geq 1}$, one can see that $x^0 \in \Phi(x^0)$. \square

4. FIXED POINT-THEOREMS FOR HOMOLOGICAL $UV^n(G)$ AND $UV^\omega(G)$ DECOMPOSITIONS

In this section we provide some fixed point-theorems for homological UV^n or homological $UV^\omega(G)$ decompositions of compact metric AR's, where G is a field. Our results are homological analogues of for homotopical UV^n and UV^ω decompositions, see [2, Theorems 3-4] and [5, Theorems 7.1-7.3]. We follow Gutev's scheme [5] of proofs applying our Theorem 1.3, Corollary 1.4 and Theorem 1.5 instead of their homotopical versions.

By a homological $UV^n(G)$ (resp., homological $UV^\omega(G)$) decomposition of a compactum X we mean an upper semi-continuous decomposition of X into compact homologically $UV^n(G)$ (resp., homologically $UV^\omega(G)$) sets. The decomposition space is denoted by X/\sim and $\pi : X \rightarrow X/\sim$ is the quotient map.

Theorem 4.1. *Let X be a compact metric AR with $\dim X \leq n$ and X/\sim be a homological $UV^{n-1}(G)$ decomposition of X . Then X/\sim has the fixed-point property.*

Proof. For any map $f : X/\sim \rightarrow X/\sim$ the set-valued map $\Phi := \pi^{-1} \circ f \circ \pi : X \rightarrow 2^X$ is usc and homologically $UV^{n-1}(G)$. Then, by Corollary 1.4, $x_0 \in \Phi(x_0)$ for some $x_0 \in X$. Hence, $f(\pi(x_0)) = \pi(x_0)$. \square

Theorem 4.2. *Let X be a compact metric AR and X/\sim be a homological $UV^{n-1}(G)$ decomposition of X with $\dim X/\sim \leq n$. Then X/\sim has the fixed-point property.*

Proof. For any map $f : X/\sim \rightarrow X/\sim$ consider the set-valued map $\Phi := \pi^{-1} \circ f : X/\sim \rightarrow 2^X$ and apply Theorem 1.3 to find a point $x_0 \in X$ with $x_0 \in \Phi(\pi(x_0))$. The last equality implies $f(\pi(x_0)) = \pi(x_0)$. \square

Theorem 4.3. *Let X be a compact metric AR and X/\sim be a homological $UV^\omega(G)$ decomposition of X . Then X/\sim has the fixed-point property.*

Proof. We repeat the proof of Theorem 4.1 using now Theorem 1.5 instead of Corollary 1.4. \square

Acknowledgments. The author would like to express his gratitude to P. Semenov and S. Bogatyĭ for their helpful comments. The author also thanks the referee for his/her careful reading and suggesting some improvements of the paper.

REFERENCES

- [1] T. Banach and R. Cauty, *A homological selection theorem implying a division theorem for Q -manifolds*, Fixed point theory and its applications, 1122, Banach Center Publ., 77, Polish Acad. Sci. Inst. Math., Warsaw (2007).
- [2] S. Bogatyĭ, *On preserving the fixed point property by mappings*, General Topology and its Relations to Modern Analysis and Algebra IV (Proc. Fourth Prague Topological Symposium, Prague (1976)), Part B, pp. 51-58.
- [3] R. Cauty, *Rétractes absolus de voisinage algébrique*, Serdica Math. J. **31**, 309-354 (2005).
- [4] L. Gorniewicz, A. Granas and W. Kryszewski, *On the homotopy method in the fixed point Index Theory of multi-valued mappings of compact absolute neighborhood retracts*, Journ. Math. Anal. Appl. **161**, 457-473 (1991).
- [5] V. Gutev, *Selections and approximations in finite-dimensional spaces*, Topol. Appl. **146-147**, 353-383 (2005).
- [6] V. Gutev and V. Valov, *Dense families of selections and finite-dimensional spaces*, Set. Val. Anal. **11**, 373-391 (2003).
- [7] V. Gutev, *A fixed point theorem for UV^n usco maps*, Proc. Am. Math. Soc. **124**(3), 945-952 (1996).
- [8] K. Sakai, *Geometric aspects of general topology*, Springer Monographs in Mathematics. Springer, Tokyo (2013).
- [9] E. Spanier, *Algebraic Topology*, McGraw-Hill Book Company (1966).
- [10] V. Uspenskij, *A selection theorem for C -compacta*, Topol. Appl. **85**, 351-374 (1998).

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. Box 5002, NORTH BAY, ON, P1B 8L7, CANADA

E-mail address: veskov@nipissingu.ca